If you have **high blood glucose**, make notes in your log and talk with your health care team about possible changes to your meal plan, physical activity, or diabetes medications.

Low blood glucose occurs when your blood glucose level drops below 70 mg/dL. Symptoms include:

- hunger
- nervousness and shakiness
- sweating
- light-headedness or confusion
- sleepiness

If you think your blood glucose is too low, check it. If it's 70 mg/dL or less, have one of these items right

- 2 to 5 glucose tablets
- 1/2 cup (4 ounces) of fruit juice
- 1/2 cup (4 ounces) of a regular (not diet) soft drink
- 8 ounces of milk
- 5 to 7 pieces of hard candy
- 2 teaspoons of sugar or honey

After 15 minutes, check your blood glucose again. If it's still below 70 mg/dL, have another serving. Repeat these steps until your blood glucose is at 70 mg/dL.

Blood Glucose Log

Name:
Date: Fromto
My Doctor
Name:
Phone:
My Diabetes Educator
Name:
Phone:
My blood glucose targets
Before meals:to
2 hours after the start of a meal: Less than

TO MAKE MORE MONTHLY LOGS: Make one (1) copy of this page and two (2) copies of the next page. Cut the pages in half, placing this page on top. Staple in the upper left-hand corner and fold to fit in your pocket or purse.

FOR DIABETES INFORMATION: Contact the American Diabetes Association at 1-800-DIABETES (342-2383) or visit diabetes.org. ©2004 by the American Diabetes Association, Inc. 03/04

---- CUT HERE ><-----

----- CUT HERE ><---

Date	Time	Breakfast	Medication/Comment	Time	Lunch	Medication/Comment	Time	Dinner	Medication/Comment	Time	Snack/Other	Medication/Comment

Date	Time	Breakfast	Medication/Comment	Time	Lunch	Medication/Comment	Time	Dinner	Medication/Comment	Time	Snack/Other	Medication/Comment

Date	Time	Breakfast	Medication/Comment	Time	Lunch	Medication/Comment	Time	Dinner	Medication/Comment	Time	Snack/Other	Medication/Comment