

***Clinical Conversations on the Diagnosis and Management of Fibromyalgia:
Focusing on Physician Assistants in Primary Care***

Unapproved Product Uses

Generic Name	Trade Name	Approved Use (if any)	Unapproved/ Investigational Use
Amitriptyline	Elavil Endep Vanatrip	<ul style="list-style-type: none"> • Treatment of major depression 	<ul style="list-style-type: none"> • Treatment of fibromyalgia
Cyclobenzaprine (sleep)	Amrix Fexmid Flexeril	<ul style="list-style-type: none"> • Adjunct to rest and physical therapy for relief of muscle spasm associated with acute, painful musculoskeletal conditions 	<ul style="list-style-type: none"> • Treatment of fibromyalgia
Fluoxetine	Prozac	<ul style="list-style-type: none"> • Treatment of major depressive disorder in adult and pediatric patients age 8–18 • Treatment of obsessive compulsive disorder in adult and pediatric patients age 7–17 • Treatment of bulimia nervosa in adults • Treatment of panic disorder, with or without agoraphobia, in adults 	<ul style="list-style-type: none"> • Treatment of fibromyalgia
Gabapentin	Neurontin	<ul style="list-style-type: none"> • Treatment of postherpetic neuralgia • Adjunctive therapy for partial seizures with or without secondary generalization in patients over the age of 12 with epilepsy. Also used as adjunctive therapy in patients with partial seizures from the age of 3–12 	<ul style="list-style-type: none"> • Treatment of fibromyalgia
Lidocaine (combined with marcaine for tender point injections)	Xylocaine Lidoderm patch	<ul style="list-style-type: none"> • Acute treatment of ventricular arrhythmias such as those occurring during acute myocardial infarction or during cardiac manipulation (Xylocaine) • Treatment of pain in patients with postherpetic neuralgia (Lidoderm) 	<ul style="list-style-type: none"> • In combination with marcaine to relieve tender point pain in patients with fibromyalgia

***Clinical Conversations on the Diagnosis and Management of Fibromyalgia:
Focusing on Physician Assistants in Primary Care***

Unapproved Product Uses

Metaxalone	Skelaxin	<ul style="list-style-type: none"> • An adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute, painful musculoskeletal conditions 	<ul style="list-style-type: none"> • Treatment of stiffness in patients with fibromyalgia
Methocarbamol (rigidity in fibro)	Robaxin	<ul style="list-style-type: none"> • An adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute, painful musculoskeletal conditions 	<ul style="list-style-type: none"> • Treatment of stiffness in patients with fibromyalgia
Modafinil (increase energy, decrease fatigue)	Provigil	<ul style="list-style-type: none"> • Improve wakefulness in patients with excessive sleepiness associated with narcolepsy, obstructive sleep apnea, and shift work disorder 	<ul style="list-style-type: none"> • Improve symptoms of fatigue in patients with fibromyalgia
Venlafaxine	Effexor	<ul style="list-style-type: none"> • Treatment of major depressive disorder, generalized anxiety disorder, social anxiety disorder, and panic disorder 	<ul style="list-style-type: none"> • Treatment of fibromyalgia